

## ON DIVERGENCES OF 6D, $\mathcal{N} = (1, 0)$ HYPERMULTIPLET SELF-COUPLING MODEL

B. S. Merzlikin<sup>1,2,\*</sup>, L. L. Ryskina<sup>3,\*\*</sup>

<sup>1</sup> Tomsk Polytechnic University, Tomsk, Russia

<sup>2</sup> Tomsk State University of Control Systems and Radioelectronics, Tomsk, Russia

<sup>3</sup> Siberian State Medical University, Tomsk, Russia

We consider the six-dimensional  $\mathcal{N} = (1, 0)$  supersymmetric hypermultiplet model with arbitrary self-coupling in the external classical gauge superfield background. We use the background field method in harmonic superspace to study the one-loop divergences of the effective action of the model.

Рассматривается шестимерная суперсимметрическая модель  $\mathcal{N} = (1, 0)$  гипермультиплета с произвольным самодействием на фоне внешнего классического калибровочного суперполя. С использованием метода фонового поля в гармоническом суперпространстве изучена структура однопетлевых расходимостей эффективного действия в модели.

PACS: 12.60.Jv

---

\* E-mail: merzlikin@tspu.edu.ru

\*\* E-mail: ryskina.ll@ssmu.ru